


1 Glueability of resource proof-structures: inverting 2 the Taylor expansion

3 Giulio Guerrieri 

4 University of Bath, Department of Computer Science, Bath, UK
5 g.guerrieri@bath.ac.uk

6 Luc Pellissier

7 Université de Paris, IRIF, CNRS, F-75013 Paris, France
8 pellissier@irif.fr

9 Lorenzo Tortora de Falco

10 Università Roma Tre, Dipartimento di Matematica e Fisica, Rome, Italy
11 tortora@uniroma3.it

12 — Abstract —

13 A Multiplicative-Exponential Linear Logic (MELL) proof-structure can be expanded into a set of
14 resource proof-structures: its Taylor expansion. We introduce a new criterion characterizing those
15 sets of resource proof-structures that are part of the Taylor expansion of some MELL proof-structure,
16 through a rewriting system acting both on resource and MELL proof-structures.

17 **2012 ACM Subject Classification** Theory of computation → Linear logic

18 **Keywords and phrases** linear logic, Taylor expansion, proof-net, natural transformation

19 **Digital Object Identifier** 10.4230/LIPIcs.CSL.2020.24

20 **Related Version** A full version [?] of the paper is available at <http://arxiv.org/abs/1910.07936>.

21 **1** Introduction

22 **Resource λ -calculus and the Taylor expansion** Girard’s linear logic (LL, [15]) is a refine-
23 ment of intuitionistic and classical logic that isolates the infinitary parts of reasoning in two
24 (dual) modalities: the *exponentials* ! and ?. They give a logical status to the operations of
25 memory management such as *copying* and *erasing*: a linear proof corresponds—via Curry–
26 Howard isomorphism—to a program that uses its argument *linearly*, *i.e.* exactly once, while
27 an exponential proof corresponds to a program that can use its argument at will.

28 The intuition that linear programs are analogous to linear functions (as studied in linear
29 algebra) while exponential programs mirror a more general class of analytic functions got a
30 technical incarnation in Ehrhard’s work [9, 10] on LL-based denotational semantics for the
31 λ -calculus. This investigation has been then internalized in the syntax, yielding the *resource*
32 *λ -calculus* [5, 11, 14]: there, copying and erasing are forbidden and replaced by the possibility
33 to apply a function to a *bag* of resource λ -terms which specifies how many times an argument
34 can be linearly passed to the function, so as to represent only bounded computations.

35 The *Taylor expansion* associates with an ordinary λ -term a (generally infinite) set of
36 resource λ -terms, recursively approximating the usual application: the Taylor expansion of
37 the λ -term MN is made of resource λ -terms of the form $t[u_1, \dots, u_n]$, where t is a resource
38 λ -term in the Taylor expansions of M , and $[u_1, \dots, u_n]$ is a bag of arbitrarily finitely many
39 (possibly 0) resource λ -terms in the Taylor expansion of N . Roughly, the idea is to decompose
40 a program into a set of purely “resource-sensitive programs”, all of them containing only
41 bounded (although possibly non-linear) calls to inputs. The notion of Taylor expansion has
42 many applications in the theory of the λ -calculus, *e.g.* in the study of linear head reduction
43 [12], normalization [23, 26], Böhm trees [4, 18], λ -theories [19], intersection types [21]. More



© Giulio Guerrieri and Luc Pellissier and Lorenzo Tortora de Falco;
licensed under Creative Commons License CC-BY

28th EACSL Annual Conference on Computer Science Logic (CSL 2020).

Editors: Maribel Fernández and Anca Muscholl; Article No. 24; pp. 24:1–24:17



Leibniz International Proceedings in Informatics

Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

44 generally, understanding the relation between a program and its Taylor expansion renews the
 45 logical approach to the quantitative analysis of computation started with the inception of LL.

46 A natural question is the *inverse Taylor expansion problem*: how to characterize which
 47 sets of resource λ -terms are contained in the Taylor expansion of a same λ -term? Ehrhard and
 48 Regnier [14] defined a simple *coherence* relation such that a finite set of resource λ -terms is
 49 included in the Taylor expansion of a λ -term if and only if the elements of this set are pairwise
 50 coherent. Coherence is crucial in many structural properties of the resource λ -calculus, such
 51 as in the proof that in the λ -calculus normalization and Taylor expansion commute [12, 14].

52 We aim to solve the inverse Taylor expansion problem in the more general context of LL,
 53 more precisely in the *multiplicative-exponential fragment* MELL of LL, being aware that for
 54 MELL no coherence relation can solve the problem (see below).

55 **Proof-nets, proof-structures and their Taylor expansion: seeing trees behind graphs** In
 56 MELL, linearity and the sharp analysis of computations naturally lead to represent proofs
 57 in a more general *graph*-like syntax instead of a term-like or tree-like one.¹ Indeed, linear
 58 negation is involutive and classical duality can be interpreted as the possibility of juggling
 59 between different conclusions, without a distinguished output. Graphs representing proofs in
 60 MELL are called *proof-nets*: their syntax is richer and more expressive than the λ -calculus.
 61 Contrary to λ -terms, proof-nets are special inhabitants of the wider land of *proof-structures*:
 62 they can be characterized, among proof-structures, by abstract (geometric) conditions called
 63 correctness criteria [15]. The procedure of cut-elimination can be applied to proof-structures,
 64 and proof-nets can also be seen as the proof-structures with a good behavior with respect to
 65 cut-elimination [1]. Proof-structures can be interpreted in denotational models and proof-
 66 nets can be characterized among them by semantic means [24]. It is then natural to attack
 67 problems in the general framework of proof-structures. In this work, correctness plays no role
 68 at all, hence we will consider proof-structures and not only proof-nets. MELL proof-structures
 69 are a particular kind of graphs, whose edges are labeled by MELL formulæ and vertices by
 70 MELL connectives, and for which special subgraphs are highlighted, the *boxes*, representing
 71 the parts of the proof-structure that can be copied and discarded (*i.e.* called an unbounded
 72 number of times). A box is delimited from the rest of a proof-structure by exponential
 73 modalities: its border is made of one !-cell, its principal door, and arbitrarily many ?-cells,
 74 its auxiliary doors. Boxes are nested or disjoint (they cannot partially overlap), so as to add
 75 a tree-like structure to proof-structures *aside* from their graph-like nature.

76 As in λ -calculus, one can define [13] box-free *resource proof-structures*², where !-cells make
 77 resources available boundedly, and the *Taylor expansion* of MELL proof-structures into these
 78 resource proof-structures, that recursively copies the content of the boxes an arbitrary number
 79 of times. In fact, as somehow anticipated by Boudes [3], such a Taylor expansion operation can
 80 be carried on any tree-like structure. This primitive, abstract, notion of Taylor expansion can
 81 then be pulled back to the structure of interest, as shown in [17] and put forth again here.

82 **The question of coherence for proof-structures** The inverse Taylor expansion problem
 83 has a natural counterpart in the world of MELL proof-structures: given a set of resource
 84 proof-structures, is there a MELL proof-structure the expansion of which contains the set?
 85 Pagani and Tasson [22] give the following answer: it is possible to decide whether a finite set of
 86 resource proof-structures is a subset of the Taylor expansion of a same MELL proof-structure

¹ A term-like object is essentially a tree, with one output (its root) and many inputs (its other leaves).

² Also known as differential proof-structures [6] or differential nets [13, 20, 7] or simple nets [22].

87 (and even possible to do it in non-deterministic polynomial time); but unlike the λ -calculus,
 88 the structure of the relation “being part of the Taylor expansion of a same proof-structure”
 89 is *much more* complicated than a binary (or even n -ary) coherence. Indeed, for any $n > 1$, it
 90 is possible to find $n + 1$ resource proof-structures such that any n of them are in the Taylor
 91 expansion of some MELL proof-structure, but there is no MELL proof-structure whose Taylor
 92 expansion has all the $n + 1$ as elements (see our Example 21 and [25, pp. 244-246]).


93 In this work, we introduce a new combinatorial criterion, *glueability*, for deciding whether
 94 a set of resource proof-structures is a subset of the Taylor expansion of some MELL proof
 95 structure, based on a rewriting system on sequences of MELL formulæ. Our criterion is more
 96 general (and, we believe, simpler) than the one of [22], which is limited to the *cut-free* case with
 97 *atomic axioms* and characterizes only *finite* sets: we do not have these limitations. We believe
 98 that our criterion is a useful tool for studying proof-structures. We conjecture that it can be
 99 used to show that, for a suitable geometric restriction, a binary coherence relation does exist
 100 for resource proof-structures. It might also shed light on correctness and sequentialization.

101 As the proof-structures we consider are typed, an unrelated difficulty arises: a resource
 102 proof-structure might not be in the Taylor expansion of any MELL proof-structure, not
 103 because it does not respect the structure imposed by the Taylor expansion, but because its
 104 type is impossible.³ To solve this issue we enrich the MELL proof-structure syntax with a
 105 “universal” proof-structure: a special \boxtimes -cell (*daimon*) that can have any number of outputs
 106 of any types, and we allow it to appear inside a box, representing information plainly missing
 107 (see Section 8 for more details and the way this matter is handled by Pagani and Tasson [22]).

108 2 Outline and technical issues

109 **The rewritings** The essence of our rewriting system is not located on proof-structures but
 110 on lists of MELL formulæ (Definition 9). In a very down-to-earth way, this rewriting system is
 111 generated by elementary steps akin to rules of sequent calculus read from the *bottom up*: they
 112 act on a list of conclusions, analogous to a monolaterous right-handed sequent. These steps are
 113 actually more sequentialized than sequent calculus rules, as they do not allow for commutation.
 114 For instance, the rule corresponding to the introduction of a \otimes on the i -th formula, is defined
 115 as $\otimes_i : (\gamma_1, \dots, \gamma_{i-1}, A \otimes B, \gamma_{i+1}, \dots, \gamma_n) \rightarrow (\gamma_1, \dots, \gamma_{i-1}, A, B, \gamma_{i+1}, \dots, \gamma_n)$.

116 These rewrite steps then act on MELL proof-structures, coherently
 117 with their type, by modifying (most of the times, erasing) the cells
 118 directly connected to the conclusion of the proof-structure. Formally,
 119 this means that there is a functor $\mathbf{qMELL}^{\boxtimes}$ from the rewrite steps



120 into the category \mathbf{Rel} of sets and relations, associating with a list of formulæ the set of MELL
 121 proof-structures with these conclusions, and with a rewrite step a relation implementing it
 122 (Definition 12). The rules *deconstruct* the proof-structure, starting from its conclusions. The
 123 rule \otimes_1 acts by removing a \otimes -cell on the first conclusion, replacing it by two conclusions.

124 These rules can only act on specific proof-structures, and indeed, capture a lot of their
 125 structure: \otimes_i can be applied to a MELL proof-structure R if and only if R has a \otimes -cell in
 126 the conclusion i (as opposed to, say, an axiom). So, in particular, every proof-structure is
 127 completely characterized by any sequence rewriting it to the empty proof-structure.

³ Similarly, in the λ -calculus, there is no closed λ -term of type $X \rightarrow Y$ with $X \neq Y$ atomic, but the resource λ -term $(\lambda f.f)[\]$ can be given that type: the empty bag $[\]$ kills any information on the argument.


24:4 Glueability of resource proof-structures: inverting the Taylor expansion

128 **Naturality** The same rules act also on sets of resource proof-structures, defining the functor
 129 $\mathfrak{PqDiLL}_0^{\mathfrak{X}}$ from the rewrite steps into the category **Rel** (Definition 17). When carefully
 130 defined, the Taylor expansion induces a *natural transformation* from $\mathfrak{PqDiLL}_0^{\mathfrak{X}}$ to $\mathfrak{qMELL}^{\mathfrak{X}}$
 131 (Theorem 18). By applying this naturality repeatedly, we get our characterization (The-
 132 orem 20): a set of resource proof-structures Π is a subset of the Taylor expansion of a MELL
 133 proof-structure iff there is a sequence rewriting Π to the singleton of the *empty* proof-structure.

134 The naturality property is not only a mean to get our characterization, but also an
 135 interesting result in itself: natural transformations can often be used to express fundamental
 136 properties in a mathematical context. In this case, the *Taylor expansion is natural* with
 137 respect to the possibility to build a proof-structure (both MELL or resource) by adding a cell
 138 to its conclusions or boxing it. Said differently, naturality of the Taylor expansion roughly
 139 means that the rewrite rules that deconstruct a MELL proof-structure R and a set of resource
 140 proof-structures in the Taylor expansion of R mimic each other.

141 **Quasi-proof-structures and mix** Our rewrite rules consume proof-structures from their
 142 conclusions. The rule corresponding to boxes in MELL opens a box by deleting its principal
 143 door (a !-cell) and its border, while for a resource proof-structure it deletes a !-cell and
 144 separates the different copies of the content of the box (possibly) represented by such a !-cell.
 145 This operation is problematic in a twofold way. In a resource proof-structure, where the
 146 border of boxes is not marked, it is not clear how to identify such copies. On the other side,
 147 in a MELL proof-structure the content of a box is not to be treated as if it were at the same
 148 level as what is outside of the box: it can be copied many times or erased, while what is
 149 outside boxes cannot, and treating the content in the same way as the outside suppresses
 150 this distinction, which is crucial in LL. So, we need to remember that the content of a box,
 151 even if it is at depth 0 (*i.e.* not contained in any other box) after erasing the box wrapping
 152 it by means of our rewrite rules, is not to be mixed with the rest of the structure at depth 0.


153 In order for our proof-structures to provide this information, we need to
 154 generalize them and consider that a proof-structure can have not just a tree of
 155 boxes, but a *forest*: this yields the notion of *quasi-proof-structure* (Definition 1).



156 In this way, according to our rewrite rules, opening a box by deleting its principal door
 157 amounts to taking a box in the tree and disconnecting it from its root, creating a new tree.
 158 We draw this by surrounding elements having the same root with a dashed line, open from
 159 the bottom, remembering the phantom presence of the border of the box, below, even if it
 160 was erased. This allows one to open the box only when it is “alone” (see Definition 11).

161 This is not merely a technical remark, as this generalization gives a status to the *mix*
 162 rule of LL: indeed, mixing two proofs amounts to taking two proofs and considering them
 163 as one, without any other modifications. Here, it amounts to taking two proofs, each with
 164 its box-tree, and considering them as one by merging the roots of their trees (see the *mix*
 165 step in Definition 11). We embed this design decision up to the level of formulæ, which
 166 are segregated in different zones that have to be mixed before interacting (see the notion of
 167 partition of a finite sequence of formulæ in Section 3).

168 **Geometric invariance and emptiness: the filled Taylor expansion** The use of forests
 169 instead of trees for the nesting structure of boxes, where the different roots are thought of
 170 as the contents of long-gone boxes, has an interesting consequence in the Taylor expansion:
 171 indeed, an element of the Taylor expansion of a proof-structure contains an arbitrary number
 172 of copies of the contents of the boxes, in particular *zero*. If we think of the part at depth
 173 0 of a MELL proof-structure as inside an invisible box, its content can be deleted in some



■ **Figure 1** Cells, with their labels and their typed inputs and outputs (ordered from left to right).

174 elements of the Taylor expansion just as any other box.⁴ As erasing completely conclusions
 175 would cause the Taylor expansion not preserve the conclusions (which would lead to technical
 176 complications), we introduce the *filled Taylor expansion* (Definition 8), which contains not
 177 only the elements of the usual Taylor expansion, but also elements of the Taylor expansion
 178 where one component has been erased and replaced by a \boxtimes -cell (*daimon*), representing a
 179 lack of information, apart from the number and types of the conclusions.

180 **Atomic axioms** Our paper first focuses on the case where proof-structures are restricted to
 181 *atomic axioms*. In Section 7 we sketch how to adapt our method to the non-atomic case.

182 3 Proof-structures and the Taylor expansion

183 **MELL formulæ and (quasi-)proof-structures** Given a countably infinite set of propositional
 184 variables X, Y, Z, \dots , MELL *formulæ* are defined by the following inductive grammar:

$$185 \quad A, B ::= X \mid X^\perp \mid \mathbf{1} \mid \perp \mid A \otimes B \mid A \wp B \mid !A \mid ?A$$

186 Linear negation is defined via De Morgan laws $\mathbf{1}^\perp = \perp$, $(A \otimes B)^\perp = A^\perp \wp B^\perp$ and
 187 $(!A)^\perp = ?A$, so as to be involutive, *i.e.* $A^{\perp\perp} = A$. Given a list $\Gamma = (A_1, \dots, A_m)$ of MELL
 188 formulæ, a *partition* of Γ is a list $(\Gamma_1, \dots, \Gamma_n)$ of lists of MELL formulæ such that there are
 189 $0 = i_0 < \dots < i_n = m$ with $\Gamma_j = (A_{i_{j-1}+1}, \dots, A_{i_j})$ for all $1 \leq j \leq n$; such a partition of Γ
 190 is also denoted by $(A_1, \dots, A_{i_1}; \dots; A_{i_{n-1}+1}, \dots, A_m)$, with lists separated by semi-colons.

191 We reuse the syntax of proof-structures given in [17] and sketch here its main features. We
 192 suppose known definitions of (directed) graph, rooted tree, and morphism of these structures.
 193 In what follows we will speak of *tails* in a graph: “hanging” edges with only one vertex. This
 194 can be implemented either by adding special vertices or using [2]’s graphs.

195 If an edge e is incoming in (resp. outgoing from) a vertex v , we say that e is a *input*
 196 (resp. *output*) of v . The reflexive-transitive closure of a tree τ is denoted by τ° : the operator
 197 $(\cdot)^\circ$ lifts to a functor from the category of trees to the category of directed graphs.

198 ► **Definition 1.** A module M is a (finite) directed graph with:

- 199 ■ vertices v labeled by $\ell(v) \in \{\mathbf{ax}, \mathbf{cut}, \mathbf{1}, \perp, \otimes, \wp, ?, !\} \cup \{\boxtimes_p \mid p \in \mathbb{N}\}$, the type of v ;
 - 200 ■ edges e labeled by a MELL formula $\mathbf{c}(e)$, the type of e ;
 - 201 ■ an order $<_M$ that is total on the tails of $|M|$ and on the inputs of each vertex of type \wp, \otimes .
- 202 Moreover, all the vertices verify the conditions of Figure 1.⁵

203 A quasi-proof-structure is a triple $R = (|R|, \mathcal{F}, \mathbf{box})$ where:

- 204 ■ $|R|$ is a module with no input tails, called the module of R ;
- 205 ■ \mathcal{F} is a forest of rooted trees with no input tails, called the box-forest of R ;
- 206 ■ $\mathbf{box}: |R| \rightarrow \mathcal{F}^\circ$ is a morphism of directed graphs, the box-function of R , which induces a
 207 partial bijection from the inputs of the vertices of type $!$ and the edges in \mathcal{F} , and such that:

⁴ The dual case, of copying the contents of a box, poses no problem in our approach.

⁵ Note that there are no conditions on the types of the outputs of vertices of type \boxtimes (*i.e.* of type \boxtimes_p for some $p \in \mathbb{N}$); and the outputs of vertices of type \mathbf{ax} must have *atomic* types.

208 ■ for any vertices v, v' with an edge from v' to v , if $\text{box}(v) \neq \text{box}(v')$ then $\ell(v) \in \{!, ?\}$.⁶
 209 Moreover, for any output tails e_1, e_2, e_3 in $|R|$ which are outputs of the vertices v_1, v_2, v_3 ,
 210 respectively, if $e_1 <_{|R|} e_2 <_{|R|} e_3$ then it is impossible that $\text{box}(v_1) = \text{box}(v_3) \neq \text{box}(v_2)$.⁷

211 A quasi-proof-structure $R = (|R|, \mathcal{F}, \text{box})$ is:

- 212 1. MELL^{\boxtimes} if all vertices in $|R|$ of type $!$ have exactly one input, and the partial bijection
- 213 induced by box from the inputs of the vertices of type $!$ in $|R|$ and the edges in \mathcal{F} is total.
- 214 2. MELL if it is MELL^{\boxtimes} and, for every vertex v in $|R|$ of type \boxtimes , one has $\text{box}^{-1}(\text{box}(v)) = \{v\}$
- 215 and $\text{box}(v)$ is not a root of the box-forest \mathcal{F} of R .
- 216 3. $\text{DiLL}_0^{\boxtimes}$ if the box-forest \mathcal{F} of R is just a juxtaposition of roots.
- 217 4. DiLL_0 (or resource) if it is $\text{DiLL}_0^{\boxtimes}$ and there is no vertex in $|R|$ of type \boxtimes .

218 For the previous systems, a proof-structure is a quasi-proof-structure whose box-forest is a tree.

219 Our MELL proof-structure (i.e. a MELL quasi-proof-structure that is also a proof-structure)
 220 corresponds to the usual notion of MELL proof-structure (as in [8]) except that we also allow
 221 the presence of a box filled only by a *daimon* (i.e. a vertex of type \boxtimes). The *empty* (DiLL_0 and
 222 MELL) proof-structure—whose module and box-forest are empty graphs—is denoted by ε .


223 Given a quasi-proof-structure $R = (|R|, \mathcal{F}, \text{box})$, the output tails of $|R|$ are the *conclusions*
 224 of R . So, the pre-images of the roots of \mathcal{F} via box partition the conclusions of R in a list of
 225 lists of such conclusions. The *type* of R is the list of lists of the types of these conclusions.
 226 We often identify the conclusions of R with a finite initial segment of \mathbb{N} .

227 By definition of graph morphism, two conclusions in two distinct lists in the type of a
 228 quasi-proof-structure R are in two distinct connected components of $|R|$; so, if R is not a
 229 proof-structure then $|R|$ contains several connected components. Thus, R can be seen as a
 230 list of proof-structures, its *components*, one for each root in its box-forest.

231 A non-root vertex v in the box-forest \mathcal{F} induces a subgraph of \mathcal{F}° of all vertices above it
 232 and edges connecting them. The pre-image of this subgraph through box is the *box* of v and
 233 the conditions on box in Definition 1 translate the usual nesting condition for LL boxes.

234 In quasi-proof-structures, we speak of *cells* instead of vertices, and, for a cell of type ℓ , of
 235 a ℓ -cell. A \boxtimes -cell is a \boxtimes_p -cell for some $p \in \mathbb{N}$. An *hypothesis cell* is a cell without inputs.

236 ► **Example 2.** The graph in Figure 2 is a MELL quasi-proof-structure. The colored areas
 237 represent the pre-images of boxes, and the dashed boxes represent the pre-images of roots.



■ **Figure 2** A MELL quasi-proof-structure R , its box-forest \mathcal{F}_R (without dotted lines) and the reflexive-transitive closure \mathcal{F}_R° of \mathcal{F}_R (with also dotted lines).


⁶ Roughly, it says that the border of a box is made of (inputs of) vertices of type $!$ or $?$.

⁷ This is a technical condition that simplifies the definition of the rewrite rules in Section 4. Note that $\text{box}(v_1), \text{box}(v_2), \text{box}(v_3)$ are necessarily roots in \mathcal{F} , since box is a morphism of directed graphs.

238 **The Taylor expansion** Proof-structures have a tree structure made explicit by their box-
 239 function. Following [17], the definition of the Taylor expansion uses this tree structure: first,
 240 we define how to “*expand*” a tree—and more generally a forest—via a generalization of the
 241 notion of thick subtree [3] (Definition 3; roughly, a thick subforest of a box-forest says the
 242 number of copies of each box to be taken, iteratively), we then take all the expansions of the
 243 tree structure of a proof-structure and we *pull* the approximations *back* to the underlying
 244 graphs (Definition 5), finally we *forget* the tree structures associated with them (Definition 6).

245 ► **Definition 3** (thick subforest). *Let τ be a forest of rooted trees. A thick subforest of τ is a*
 246 *pair (σ, h) of a forest σ of rooted trees and a graph morphism $h: \sigma \rightarrow \tau$ whose restriction to*
 247 *the roots of σ is bijective.*

248 ► **Example 4.** The following is a graphical presentation of a thick subforest (τ, h) of the
 249 box-forest \mathcal{F} of the quasi-proof-structure in Figure 2, where the graph morphism $h: \tau \rightarrow \mathcal{F}$
 250 is depicted chromatically (same color means same image via h).



251
 252
 253 Intuitively, it means that τ is obtained from \mathcal{F} by taking 3 copies of the blue box, 1 copy of
 254 the red box and 4 copies of the orange box; in the first (resp. second; third) copy of the blue
 255 box, 1 copy (resp. 0 copies; 2 copies) of the purple box has been taken.

256 ► **Definition 5** (proto-Taylor expansion). *Let $R = (|R|, \mathcal{F}_R, \text{box}_R)$ be a quasi-proof-structure.*
 257 *The proto-Taylor expansion of R is the set $\mathcal{T}^{\text{proto}}(R)$ of thick subforests of \mathcal{F}_R .*

258 *Let $t = (\tau_t, h_t) \in \mathcal{T}^{\text{proto}}(R)$. The t -expansion of R is the pullback (R_t, p_t, p_R) below,*
 259 *computed in the category of directed graphs and graph morphisms.*

$$\begin{array}{ccc}
 R_t & \xrightarrow{p_t} & \tau_t^\circ \\
 \downarrow p_R & \lrcorner & \downarrow h_t^\circ \\
 |R| & \xrightarrow{\text{box}_R} & \mathcal{F}_R^\circ
 \end{array}$$


260

261 Given a quasi-proof-structure R and $t = (\tau_t, h_t) \in \mathcal{T}^{\text{proto}}(R)$, the directed graph R_t
 262 inherits labels on vertices and edges by composition with the graph morphism $p_R: R_t \rightarrow |R|$.

263 Let $[\tau_t]$ be the forest made up of the roots of τ_t and $\iota: \tau_t \rightarrow [\tau_t]$ be the graph morphism
 264 sending each vertex of τ_t to the root below it; ι° induces by post-composition a morphism
 265 $\bar{h}_t = \iota^\circ \circ p_t: R_t \rightarrow [\tau_t]^\circ$. The triple $(R_t, [\tau_t], \bar{h}_t)$ is a DiLL_0 quasi-proof-structure, and it is a
 266 DiLL_0 proof-structure if R is a proof-structure. We can then define the *Taylor expansion* $\mathcal{T}(R)$
 267 of a quasi-proof-structure R (an example of an element of a Taylor expansion is in Figure 3).

268 ► **Definition 6** (Taylor expansion). *Let R be a quasi-proof-structure. The Taylor expansion of*
 269 *R is the set of DiLL_0 quasi-proof-structures $\mathcal{T}(R) = \{(R_t, [\tau_t], \bar{h}_t) \mid t = (\tau_t, h_t) \in \mathcal{T}^{\text{proto}}(R)\}$.*

270 An element $(R_t, [\tau_t], \bar{h}_t)$ of the Taylor expansion of a quasi-proof-structure R has much
 271 less structure than the pullback (R_t, p_t, p_R) : the latter indeed is a DiLL_0 quasi-proof-structure
 272 R_t coming with its projections $|R| \xleftarrow{p_R} R_t \xrightarrow{p_t} \tau_t^\circ$, which establish a precise correspondence
 273 between cells and edges of R_t and cells and edges of R : a cell in R_t is labeled (via the
 274 projections) by both the cell of $|R|$ and the branch of the box-forest of R it arose from. But
 275 $(R_t, [\tau_t], \bar{h}_t)$ where R_t is without its projections p_t and p_R loses the correspondence with R .




■ **Figure 3** The element of the Taylor expansion of the MELL quasi-proof-structure R in Figure 2, obtained from the element of $\mathcal{T}^{\text{proto}}(R)$ depicted in Example 4.

276 ▶ **Remark 7.** By definition, the Taylor expansion preserves conclusions: there is a bijection
 277 φ from the conclusions of a quasi-proof-structure R to the ones in each element ρ of $\mathcal{T}(R)$
 278 such that i and $\varphi(i)$ have the same type and the same root (i.e. $\text{box}_R(i) = \text{box}_\rho(\varphi(i))$ up to
 279 isomorphism). Therefore, the types of R and ρ are the same (as a list of lists).

280 **The filled Taylor expansion** As discussed in Section 2 (p. 4), our method needs to “represent”
 281 the emptiness introduced by the Taylor expansion (taking 0 copies of a box) so as to preserve
 282 the conclusions. So, an element of the *filled Taylor expansion* $\mathcal{T}^{\boxtimes}(R)$ of a quasi-proof-structure
 283 R (an example is in Figure 4) is obtained from an element of $\mathcal{T}(R)$ where a whole component
 284 can be erased and replaced by a \boxtimes -cell with the same conclusions (hence $\mathcal{T}(R) \subseteq \mathcal{T}^{\boxtimes}(R)$).

285 ▶ **Definition 8** (filled Taylor expansion). An emptying of a DiLL_0 quasi-proof-structure $\rho =$
 286 $(|\rho|, \mathcal{F}, \text{box})$ is the DiLL_0 quasi-proof-structure with the same conclusions as ρ , obtained from ρ
 287 by replacing each of the components of some roots of \mathcal{F} with a \boxtimes -cell whose outputs are tails.
 288 The filled Taylor expansion $\mathcal{T}^{\boxtimes}(R)$ of a quasi-proof-structure R is the set of all the
 289 emptyings of every element of its Taylor expansion $\mathcal{T}(R)$.



■ **Figure 4** An element of the filled Taylor expansion of the MELL quasi-proof-structure in Figure 2.

290 4 Means of destruction: unwinding MELL quasi-proof-structures

291 Our aim is to deconstruct proof-structures (be they MELL^{\boxtimes} or DiLL_0) from their conclusions.
 292 To do that, we introduce a category of rules of deconstruction. The morphisms of this category
 293 are sequences of deconstructing rules, acting on lists of lists of formulæ. These morphisms
 294 act through functors on quasi-proof-structures, exhibiting their sequential structure.

295 ▶ **Definition 9** (the category **Path**). Let **Path** be the category whose
 296 ■ objects are lists $\Gamma = (\Gamma_1; \dots; \Gamma_n)$ of lists of MELL formulæ;
 297 ■ arrows are freely generated by the elementary paths in Figure 5.
 298 We call a path any arrow $\xi: \Gamma \rightarrow \Gamma'$. We write the composition of paths without symbols and
 299 in the diagrammatic order, so, if $\xi: \Gamma \rightarrow \Gamma'$ and $\xi': \Gamma' \rightarrow \Gamma''$, $\xi\xi': \Gamma \rightarrow \Gamma''$.

$$\begin{array}{l}
(\Gamma_1; \dots; \Gamma_k, c(i), c(i+1), \Gamma'_k; \dots; \Gamma_n) \xrightarrow{\text{exc}_i} (\Gamma_1; \dots; \Gamma_k, c(i+1), c(i), \Gamma'_k; \dots; \Gamma_n) \\
(\Gamma_1; \dots; \Gamma_k, c(i), c(i+1), \Gamma'_k; \dots; \Gamma_n) \xrightarrow{\text{mix}_i} (\Gamma_1; \dots; \Gamma_k, c(i); c(i+1), \Gamma'_k; \dots; \Gamma_n) \\
(\Gamma_1; \dots; \Gamma_k; c(i), c(i+1); \Gamma_{k+2}; \dots; \Gamma_n) \xrightarrow{\text{ax}_i} (\Gamma_1; \dots; \Gamma_k; \Gamma_{k+2}; \dots; \Gamma_n) \quad \text{with } c(i) = A = c(i+1)^\perp \\
(\Gamma_1; \dots; \Gamma_k; \dots; \Gamma_n) \xrightarrow{\text{cut}_i^\perp} (\Gamma_1; \dots; \Gamma_k, c(i), c(i+1); \dots; \Gamma_n) \quad \text{with } c(i) = A = c(i+1)^\perp \\
(\Gamma_1; \dots; \Gamma_k; \Gamma_{k+1}, c(i); \Gamma_{k+2}; \dots; \Gamma_n) \xrightarrow{\mathbf{x}_i} (\Gamma_1; \dots; \Gamma_k; \Gamma_{k+2}; \dots; \Gamma_n) \\
(\Gamma_1; \dots; \Gamma_k; c(i); \Gamma_{k+2}; \dots; \Gamma_n) \xrightarrow{\mathbf{1}_i} (\Gamma_1; \dots; \Gamma_k; \Gamma_{k+2}; \dots; \Gamma_n) \quad \text{with } c(i) = \mathbf{1} \\
(\Gamma_1; \dots; \Gamma_k; c(i); \Gamma_{k+2}; \dots; \Gamma_n) \xrightarrow{\perp_i} (\Gamma_1; \dots; \Gamma_k; \Gamma_{k+2}; \dots; \Gamma_n) \quad \text{with } c(i) = \perp \\
(\Gamma_1; \dots; \Gamma_k, c(i); \dots; \Gamma_n) \xrightarrow{\otimes_i} (\Gamma_1; \dots; \Gamma_k, A, B; \dots; \Gamma_n) \quad \text{with } c(i) = A \otimes B \\
(\Gamma_1; \dots; \Gamma_k, c(i); \dots; \Gamma_n) \xrightarrow{\wp_i} (\Gamma_1; \dots; \Gamma_k, A, B; \dots; \Gamma_n) \quad \text{with } c(i) = A \wp B \\
(\Gamma_1; \dots; \Gamma_k, c(i); \dots; \Gamma_n) \xrightarrow{?_i} (\Gamma_1; \dots; \Gamma_k, ?A, ?A; \dots, \Gamma_n) \quad \text{with } c(i) = ?A \\
(\Gamma_1; \dots; \Gamma_k, c(i); \dots; \Gamma_n) \xrightarrow{?_i} (\Gamma_1; \dots; \Gamma_k, A; \dots; \Gamma_n) \quad \text{with } c(i) = ?A \\
(\Gamma_1; \dots; \Gamma_k; c(i); \Gamma_{k+2}; \dots; \Gamma_n) \xrightarrow{?_i} (\Gamma_1; \dots; \Gamma_k; \Gamma_{k+2}; \dots; \Gamma_n) \quad \text{with } c(i) = ?A \\
(\Gamma_1; \dots; ?\Gamma_k, c(i); \dots; \Gamma_n) \xrightarrow{\text{Box}_i} (\Gamma_1; \dots; ?\Gamma_k, A; \dots; \Gamma_n) \quad \text{with } c(i) = !A
\end{array}$$

■ **Figure 5** The generators of **Path**. In the source $\Gamma = (A_1, \dots, A_{i_1}; \dots; A_{i_{m-1}+1}, \dots, A_{i_n})$ of each arrow, $c(i)$ denotes the i^{th} formula in the flattening $(A_1, \dots, A_{i_1}, \dots, A_{i_{m-1}+1}, \dots, A_{i_n})$ of Γ .

300 ▶ **Example 10.** $\wp_1 \wp_2 \wp_3 \otimes_1 \otimes_3 \text{exc}_1 \text{exc}_2 \text{mix}_2 \text{ax}_1 \text{exc}_2 \text{mix}_2 \text{ax}_1 \text{ax}_1$ is a path of type
301 $((X \otimes Y^\perp) \wp ((Y \otimes Z^\perp) \wp (X^\perp \wp Z))) \longrightarrow \varepsilon$, where ε is the *empty list* of lists of formulæ.

302 We will tend to forget about exchanges and perform them silently (as it is customary, for
303 instance, in most presentations of sequent calculi).

304 The category **Path** acts on $\text{MELL}^{\mathbf{x}}$ quasi-proof-structures, exhibiting a sequential struc-
305 ture in their construction. For Γ a list of list of MELL formulæ, $\mathbf{qMELL}^{\mathbf{x}}(\Gamma)$ is the set of
306 $\text{MELL}^{\mathbf{x}}$ quasi-proof-structures of type Γ . To ease the reading of the rewrite rules acting on a
307 $\text{MELL}^{\mathbf{x}}$ quasi-proof-structures R , we will only draw the parts of R belonging to the relevant
308 component; *e.g.*, if we are interested in an **ax**-cell whose outputs are the conclusions i and

309 $i+1$, and it is the only cell in a component, we will write $\overset{\text{ax}}{\underbrace{\quad}_i \quad}_{i+1}$ ignoring the rest.

310 ▶ **Definition 11** (action of paths on MELL quasi-proof-structures). *An elementary path* $a: \Gamma \rightarrow$
311 Γ' *defines a relation* $\mathfrak{a} \subseteq \mathbf{qMELL}^{\mathbf{x}}(\Gamma) \times \mathbf{qMELL}^{\mathbf{x}}(\Gamma')$ *(the action of* a *as the smallest*
312 *relation containing all the cases in Figure 6, with the following remarks:*

313 **mix** *read in reverse, a quasi-proof-structure with two components is in relation with a proof-*
314 *structure with the same module but the two roots of such components merged.*

315 **hypothesis** *if* $a \in \{\text{ax}_i, \mathbf{x}_i, \mathbf{1}_i, \perp_i, ?_w\}$, *the rules have all in common to act by deleting a cell*
316 *without inputs that is the only cell in its component. We have drawn the axiom case in*
317 *Figure 6c, the others vary only by their number of conclusions.*

318 **cut** *read in reverse, a quasi-proof-structure with two conclusions* i *and* $i+1$ *is in relation*
319 *with the quasi-proof-structure where these two conclusions are cut. This rule, from left to*
320 *right, is non-deterministic (as there are many possible cuts).*

321 **binary multiplicatives** *these rules delete a binary connective. We have only drawn the* \otimes
322 *case in Figure 6e, the* \wp *case is similar.*


323 **contraction** *splits a* $?$ -*cell with* $h+k+2$ *inputs into two* $?$ -*cells with* $h+1$ *and* $k+1$ *inputs,*
324 *respectively.*

325 **dereliction** *only applies if the* $?$ -*cell (with 1 input) does not shift a level in the box-forest.*

326 **box** *only applies if a box (and its frontier) is alone in its component.*

327 This definition of the rewrite system is extended to define a relation $\mathfrak{X} \subseteq \mathbf{qMELL}^{\mathbf{x}}(\Gamma) \times$
328 $\mathbf{qMELL}^{\mathbf{x}}(\Gamma')$ *(the action of any path* $\xi: \Gamma \rightarrow \Gamma'$ *by composition of relations.*

24:10 Glueability of resource proof-structures: inverting the Taylor expansion



■ **Figure 6** Actions of elementary paths on $\text{MELL}^{\mathfrak{X}}$ quasi-proof-structures.

329 Given two $\text{MELL}^{\mathfrak{X}}$ quasi-proof-structures R and R' , we say that a rule a applies to R if
 330 there is a finite sequence of exchanges $\text{exc}_{i_1} \cdots \text{exc}_{i_n}$ such that $R \xrightarrow{\text{exc}_{i_1} \cdots \text{exc}_{i_n} a} R'$.

331 ► **Definition 12** (the functor $\text{qMELL}^{\mathfrak{X}}$). We define a functor $\text{qMELL}^{\mathfrak{X}}: \text{Path} \rightarrow \text{Rel}$ by:
 332 ■ on objects: $\text{qMELL}^{\mathfrak{X}}(\Gamma)$ is the set of $\text{MELL}^{\mathfrak{X}}$ quasi-proof-structures of type Γ ;
 333 ■ on morphisms: for $\xi: \Gamma \rightarrow \Gamma'$, $\text{qMELL}^{\mathfrak{X}}(\xi) = \xi$ (see Definition 11).

334 Our rewrite rules enjoy two useful properties, expressed by Propositions 13 and 15.

335 ► **Proposition 13** (co-functionality). Let $\xi: \Gamma \rightarrow \Gamma'$ be a path. The relation ξ is a co-function
 336 on the sets of underlying graphs, that is, a function $\xi^{\text{op}}: \text{qMELL}^{\mathfrak{X}}(\Gamma') \rightarrow \text{qMELL}^{\mathfrak{X}}(\Gamma)$.


337 ► **Lemma 14** (applicability of rules). Let R be a non-empty $\text{MELL}^{\mathfrak{X}}$ quasi-proof-structure.
 338 There exists a conclusion i such that:

- 339 ■ either a rule in $\{\text{ax}_i, \mathbf{1}_i, \perp_i, \otimes_i, \mathfrak{X}_i, ?_i, ?_i, ?_i, \text{cut}^i, \mathfrak{X}_i, \text{Box}_i\}$ applies to R ;
- 340 ■ or $R \xrightarrow{\text{mix}_i} R'$ (where the conclusions affected by mix_i are $i-k, \dots, i, i+1, \dots, i+l$) and
 341 $i-k, \dots, i$ are all the conclusions of either a box or an hypothesis cell, and one of the
 342 components of R' coincides with this cell or box (and its border).

343 Proposition 13 and Lemma 14 are proven by simple inspection of the rewrite rules of Figure 6.

344 ► **Proposition 15** (termination). Let R be a $\text{MELL}^{\mathfrak{X}}$ quasi-proof-structure of type Γ . There
 345 exists a path $\xi: \Gamma \rightarrow \varepsilon$ such that $R \xrightarrow{\xi} \varepsilon$.

346 To prove Proposition 15, it is enough to apply Lemma 14 and show that the size of $\text{MELL}^{\mathfrak{X}}$
 347 quasi-proof-structures decreases for each application of the rules in Figure 6, according to
 348 the following definition of size. The size of a proof-structure R is the couple (p, q) where
 349 ■ p is the (finite) multiset of the number of inputs of each $?$ -cell in R ;



■ **Figure 7** Actions of elementary paths on \mathbf{X} -cells and on a box in $\mathbf{qDiLL}_0^{\mathbf{X}}$.

350 ■ q is the number of cells not labeled by \mathbf{X} in R .

351 The *size* of a quasi-proof-structure R is the (finite) multiset of the sizes of its components.

352 Multisets are ordered as usual, couples are ordered lexicographically.

353 5 Naturality of unwinding $\mathbf{DiLL}_0^{\mathbf{X}}$ quasi-proof-structures

354 For Γ a list of lists of MELL formulæ, $\mathbf{qDiLL}_0^{\mathbf{X}}(\Gamma)$ is the set of $\mathbf{DiLL}_0^{\mathbf{X}}$ quasi-proof-structures
355 of type Γ . For any set X , its powerset is denoted by $\mathfrak{P}(X)$.

356 ► **Definition 16** (action of paths on $\mathbf{DiLL}_0^{\mathbf{X}}$ quasi-proof-structures). *An elementary path*
357 *$a: \Gamma \rightarrow \Gamma'$ defines a relation $\mathfrak{a} \subseteq \mathbf{qDiLL}_0^{\mathbf{X}}(\Gamma) \times \mathfrak{P}(\mathbf{qDiLL}_0^{\mathbf{X}}(\Gamma'))$ (the action of a) by the*
358 *rules in Figure 6 (except Figure 6h, and with all the already remarked notes) and in Figure 7.*

359 *We extend this relation on $\mathfrak{P}(\mathbf{qDiLL}_0^{\mathbf{X}}(\Gamma)) \times \mathfrak{P}(\mathbf{qDiLL}_0^{\mathbf{X}}(\Gamma'))$ by the monad multiplication*
360 *of $X \mapsto \mathfrak{P}(X)$ and define \mathfrak{L} (the action of any path $\xi: \Gamma \rightarrow \Gamma'$) by composition of relations.*

361 Roughly, all the rewrite rules in Figure 7—except Figure 7h—mimic the behavior of the
362 corresponding rule in Figure 6 using a \mathbf{X} -cell. Note that in Figure 7g a \mathbf{X} -cell is created.

363 The non-empty box rule in Figure 7h requires that, on the left of \mathfrak{Box}_i , ρ_j is not connected
364 to $\rho_{j'}$ for $j \neq j'$, except for the $\mathbf{!}$ -cell and the $\mathbf{?}$ -cells in the conclusions. Read in reverse, the
365 rule associates with a non-empty finite set of \mathbf{DiLL}_0 quasi-proof-structures $\{\rho_1, \dots, \rho_n\}$ the
366 merging of ρ_1, \dots, ρ_n , that is the \mathbf{DiLL}_0 quasi-proof-structure depicted on the left of \mathfrak{Box}_i .

367 ► **Definition 17** (the functor $\mathfrak{P}\mathbf{qDiLL}_0^{\mathbf{X}}$). *We define a functor $\mathfrak{P}\mathbf{qDiLL}_0^{\mathbf{X}}: \mathbf{Path} \rightarrow \mathbf{Rel}$ by:*
368 ■ *on objects: for Γ a list of lists of MELL formulæ, $\mathfrak{P}\mathbf{qDiLL}_0^{\mathbf{X}}(\Gamma) = \mathfrak{P}(\mathbf{qDiLL}_0^{\mathbf{X}}(\Gamma))$, the*
369 *set of sets of $\mathbf{DiLL}_0^{\mathbf{X}}$ proof-structures of type Γ ;*

370 ■ *on morphisms: for $\xi: \Gamma \rightarrow \Gamma'$, $\mathfrak{P}\mathbf{qDiLL}_0^{\mathbf{X}}(\xi) = \mathfrak{L}$ (see Definition 16).*

371 ► **Theorem 18** (naturality). *The filled Taylor expansion defines a natural transformation*
372 *$\mathfrak{T}^{\mathbf{X}}: \mathfrak{P}\mathbf{qDiLL}_0^{\mathbf{X}} \Rightarrow \mathbf{qMELL}^{\mathbf{X}}: \mathbf{Path} \rightarrow \mathbf{Rel}$ by: $(\Pi, R) \in \mathfrak{T}_1^{\mathbf{X}}$ iff $\Pi \subseteq \mathcal{T}^{\mathbf{X}}(R)$ and the type of*

24:12 Glueability of resource proof-structures: inverting the Taylor expansion

373 R is Γ . Moreover, if Π is a set of DiLL_0 proof-structures with $\Pi \xrightarrow{\xi} \Pi'$ and $\Pi' \subseteq \mathcal{T}(R')$, then
 374 R is a MELL proof-structure and $\Pi \subseteq \mathcal{T}(R)$, where R is such that $R \xrightarrow{\xi} R'$.⁸

375 In other words, the following diagram commutes for every path $\xi: \Gamma \rightarrow \Gamma'$.

$$\begin{array}{ccc}
 \mathfrak{P}\mathfrak{q}\text{DiLL}_0^{\mathfrak{X}}(\Gamma) & \xrightarrow{\mathfrak{P}\mathfrak{q}\text{DiLL}_0^{\mathfrak{X}}(\xi)} & \mathfrak{P}\mathfrak{q}\text{DiLL}_0^{\mathfrak{X}}(\Gamma') \\
 \downarrow \mathfrak{T}_\Gamma^{\mathfrak{X}} & & \downarrow \mathfrak{T}_{\Gamma'}^{\mathfrak{X}} \\
 \mathfrak{q}\text{MELL}^{\mathfrak{X}}(\Gamma) & \xrightarrow{\mathfrak{q}\text{MELL}^{\mathfrak{X}}(\xi)} & \mathfrak{q}\text{MELL}^{\mathfrak{X}}(\Gamma')
 \end{array}$$

376

377 It means that given $\Pi \xrightarrow{\xi} \Pi'$, where $\Pi' \subseteq \mathcal{T}^{\mathfrak{X}}(R')$, we can simulate backwards the rewriting
 378 to R (this is where the co-functionality of the rewriting steps expressed by Proposition 13
 379 comes handy) so that $R \xrightarrow{\xi} R'$ and $\Pi \subseteq \mathcal{T}^{\mathfrak{X}}(R)$; and conversely, given $R \xrightarrow{\xi} R'$, we can
 380 simulate the rewriting for any $\Pi \subseteq \mathcal{T}^{\mathfrak{X}}(R)$, so that $\Pi \xrightarrow{\xi} \Pi'$ for some $\Pi' \subseteq \mathcal{T}^{\mathfrak{X}}(R')$.

6 Glueability of DiLL_0 quasi-proof-structures

382 Naturality (Theorem 18) allows us to characterize the sets of DiLL_0 proof-structures that are
 383 in the Taylor expansion of some MELL proof-structure (Theorem 20 below).


384 ► **Definition 19** (glueability). We say that a set Π of $\text{DiLL}_0^{\mathfrak{X}}$ quasi-proof-structures is glueable,
 385 if there exists a path ξ such that $\Pi \xrightarrow{\xi} \{\varepsilon\}$.

386 ► **Theorem 20** (glueability criterion). Let Π be a set of DiLL_0 proof-structures: Π is glueable
 387 if and only if $\Pi \subseteq \mathcal{T}(R)$ for some MELL proof-structure R .

388 **Proof.** If $\Pi \subseteq \mathcal{T}(R)$ for some MELL proof-structure R , then by termination (Proposition 15)
 389 $R \xrightarrow{\xi} \varepsilon$ for some path ξ , and so $\Pi \xrightarrow{\xi} \{\varepsilon\}$ by naturality (Theorem 18, as $\mathcal{T}^{\mathfrak{X}}(\varepsilon) = \{\varepsilon\}$).

390 Conversely, if $\Pi \xrightarrow{\xi} \{\varepsilon\}$ for some path ξ , then by naturality (Theorem 18, as $\mathcal{T}(\varepsilon) = \{\varepsilon\}$
 391 and Π is a set of DiLL_0 proof-structures) $\Pi \subseteq \mathcal{T}(R)$ for some MELL proof-structure R . ◀

392 ► **Example 21.** The three DiLL_0 proof-structures ρ_1, ρ_2, ρ_3 below are not glueable as a
 393 whole, but are glueable two by two. In fact, there is no MELL proof-structure whose Taylor
 394 expansion contains ρ_1, ρ_2, ρ_3 , but any pair of them is in the Taylor expansion of some MELL
 395 proof-structure. This is a slight variant of the example in [25, pp. 244-246].



397 An example of the action of a path starting from a DiLL_0 proof-structure ρ and ending in
 398 $\{\varepsilon\}$ can be found in Figures 8 and 9. Note that it is by no means the shortest possible path.
 399 When replayed backwards, it induces a MELL proof-structure R such that $\rho \in \mathcal{T}(R)$.

⁸ The part of the statement after “moreover” is our way to control the presence of \mathfrak{X} -cells.




Figure 8 The path $\text{Box}_2 \overset{?_1}{\rightsquigarrow} \text{Box}_2 \overset{?_2}{\rightsquigarrow} \text{mix}_1 \text{ax}_{2,3} \overset{?_1}{\rightsquigarrow} \overset{?_2}{\rightsquigarrow} \text{mix}_1 \perp_2 \overset{?_1}{\rightsquigarrow} \perp_1$ witnessing that $\rho \in \mathcal{T}(R)$ (to be continued on Figure 9).

7 Non-atomic axioms

From now on, we relax the definition of quasi-proof-structure (Definition 1 and Figure 1) so that the outputs of any ax -cell are labeled by dual MELL formulæ, not necessarily atomic. We can extend our results to this more general setting, with some technical complications. Indeed, the rewrite rule for contraction has to be modified. Consider a set of DiLL_0 proof-structures consisting of just a singleton which is a \boxtimes -cell. The contraction rule rewrites it as:


$$\begin{array}{c} \text{ax} \\ \boxtimes \\ \downarrow \downarrow \\ !A^\perp \quad !A^\perp \quad ?A \end{array} \xrightarrow{?_3} \left\{ \begin{array}{c} \text{ax} \\ \boxtimes \\ \downarrow \downarrow \\ !A^\perp \quad !A^\perp \quad ?A \quad ?A \end{array} \right\} \text{ which is then in the Taylor expansion of } \begin{array}{c} \text{ax} \\ \text{ax} \\ \downarrow \downarrow \downarrow \downarrow \\ !A^\perp \quad !A^\perp \quad ?A \quad ?A \end{array}$$

on which no contraction rewrite rule $?$ can be applied backwards, breaking the naturality. The failure of the naturality is actually due to the failure of Proposition 13 in the case of the rewrite rule $?: \overset{?}{\rightsquigarrow}^{\text{op}}$ (i.e. $\overset{?}{\rightsquigarrow}$ read from the right to the left) is functional but not total.


The solution to this conundrum lies in changing the contraction rule for $\text{DiLL}_0^{\boxtimes}$ quasi-proof-structures, by explicitly adding $?$ -cells. Hence, the application of a contraction step $?$ in the $\text{DiLL}_0^{\boxtimes}$ quasi-proof-structures precludes the possibility of anything else but a $?$ -cell on the MELL^{\boxtimes} side, which allows the contraction step $?$ to be applied backwards.

In turn, this forces us to change the definition of the filled Taylor expansion into a η -filled Taylor expansion, which has to include elements where a \boxtimes -cell (representing an empty component) has some of its outputs connected to $?$ -cells.

24:14 Glueability of resource proof-structures: inverting the Taylor expansion



■ **Figure 9** The path $\text{Box}_2 \stackrel{?}{d}_1 \text{Box}_2 \stackrel{?}{d}_2 \text{mix}_1 \text{ax}_{2,3} \stackrel{?}{c}_1 \stackrel{?}{d}_2 \text{mix}_1 \perp_2 \stackrel{?}{d}_1 \perp_1$ witnessing that $\rho \in \mathcal{T}(R)$ (continued from Figure 8).




■ **Figure 10** An element of the η -filled Taylor expansion of the MELL quasi-proof-structure in Fig. 2.

418 ► **Definition 22** (η -filled Taylor expansion). An η -emptying of a DiLL_0 quasi-proof-structure
 419 $\rho = (|\rho|, \mathcal{F}, \text{box})$ is a DiLL_0 quasi-proof-structure with the same conclusions as ρ , obtained
 420 from ρ by replacing each of the components of some roots of \mathcal{F} with a \boxtimes -cell whose outputs
 421 are either tails or inputs of a $?$ -cell whose output i is a tail, provided that i is the output tail
 422 of a $?$ -cell in ρ .

423 The η -filled Taylor expansion $\mathcal{T}_\eta^{\boxtimes}(R)$ of a quasi-proof-structure R is the set of all the
 424 η -emptyings of every element of its Taylor expansion $\mathcal{T}(R)$.

425 Note that the η -filled Taylor expansion contains all the elements of the filled Taylor
 426 expansion and some more, such as the one in Figure 10.

427 Functors $\mathbf{qMELL}^{\boxtimes}$ and $\mathfrak{P}\mathbf{qDiLL}_0^{\boxtimes}$ are defined as before (Def. 12 and 17, respectively),⁹
 428 except that the image of $\mathfrak{P}\mathbf{qDiLL}_0^{\boxtimes}$ on the generator $?_i$ (Figure 7d) is changed to



⁹ Remember that now, for Γ a list of list of MELL formulæ, $\mathbf{qMELL}^{\boxtimes}(\Gamma)$ (resp. $\mathbf{qDiLL}_0^{\boxtimes}(\Gamma)$) is the set of MELL ^{\boxtimes} (resp. DiLL ^{\boxtimes}) quasi-proof-structures of type Γ , possibly with non-atomic axioms.

431 where $?\Gamma_k$ signifies that some of the conclusions of Γ_k might be connected to the \boxtimes -cell
432 through a $?$ -cell. We can prove similarly our main results.

433 **► Theorem 23** (naturality with η). *The η -filled Taylor expansion defines a natural transform-*
434 *ation $\mathfrak{T}_\eta^\boxtimes: \mathfrak{P}\mathfrak{qDiLL}_0^\boxtimes \Rightarrow \mathfrak{qMELL}^\boxtimes: \mathbf{Path} \rightarrow \mathbf{Rel}$ by: $(\Pi, R) \in \mathfrak{T}_\eta^\boxtimes$ iff $\Pi \subseteq \mathcal{T}_\eta^\boxtimes(R)$ and the*
435 *type of R is Γ . Moreover, if Π is a set of DiLL_0 proof-structures with $\Pi \xrightarrow{\xi} \Pi'$ and $\Pi' \subseteq \mathcal{T}(R')$,*
436 *then R is a MELL proof-structure and $\Pi \subseteq \mathcal{T}(R)$, where R is such that $R \xrightarrow{\xi} R'$.*

437 **► Theorem 24** (glueability criterion with η). *Let Π be a set of DiLL_0 proof-structures, not*
438 *necessarily with atomic axioms: Π is glueable iff $\Pi \subseteq \mathcal{T}(R)$ for some MELL proof-structure R .*

439 8 Conclusions and perspectives

440 **\boxtimes -cells inside boxes** Our glueability criterion (Theorem 20) solves the inverse Taylor
441 expansion problem in a “asymmetric” way: we characterize the sets of DiLL_0 proof-structures
442 that are included in the Taylor expansion of some MELL proof-structure, but DiLL_0 proof-
443 structures have no occurrences of \boxtimes -cells, while a MELL proof-structure possibly contains
444 \boxtimes -cells inside boxes (see Definition 1). Not only this asymmetry is technically inevitable, but
445 it reflects on the fact that some glueable set of DiLL_0 proof-structure might not contain any
446 information on the content of some box (which is reified in MELL by a \boxtimes -cell), or worse that,
447 given the types, no content can fill that box. Think of the DiLL_0 proof-structure ρ made only
448 of a $!$ -cell with no inputs and one output of type $!X$, where X is atomic: $\{\rho\}$ is glueable but
449 the only MELL proof-structure R such that $\{\rho\} \subseteq \mathcal{T}(R)$ is made of a box containing a \boxtimes -cell.

450 This asymmetry is also present in Pagani and Tasson’s characterization [22], even if
451 not particularly emphasized: their Theorem 2 (analogous to the left-to-right part of our
452 Theorem 20) assumes not only that the rewriting starting from a finite set of DiLL_0 proof-
453 structures terminates but also that it ends on a MELL proof-structure (without \boxtimes -cells, which
454 ensures that there exists a MELL proof-structure without \boxtimes -cells filling all the empty boxes).

455 **The λ -calculus, connectedness and coherence** Our rewriting system and glueability cri-
456 terion should help to prove the existence of a binary coherence for elements of the Taylor
457 expansion of a fragment of MELL proof-structures (despite the impossibility for full MELL
458 proved in [25]), extending the one that exists for resource λ -terms. We can remark that our
459 glueability criterion is actually an extension of the criterion for resource λ -terms. Indeed,
460 in the case of the λ -calculus, there are three rewrite steps, corresponding to abstraction,
461 application and variable (which can be encoded in our rewrite steps), and coherence is defined
462 inductively: if a set of resource λ -terms is coherent, then any set of resource λ -term that
463 rewrites to it is also coherent.

464 Presented in this way, the main difference between the λ -calculus and MELL (concerning
465 the inverse Taylor expansion problem) would not be because of the rewriting system but
466 because the structure of any resource λ -term univocally determines the rewriting path, while,
467 for DiLL_0 proof-structures, we have to quantify existentially over all possible paths. This is
468 an unavoidable consequence of the fact that proof-structures do not have a tree-structure,
469 contrary to λ -terms and resource λ -terms.

470 Moreover, it is possible to match and mix different sequences of rewriting. Indeed,
471 consider three DiLL_0 proof-structures pairwise glueable. Proving that they are glueable as a
472 whole amounts to computing a rewriting path from the rewriting paths witnessing the three
473 glueabilities. Our paths were designed with that mixing-and-matching operation in mind, in
474 the particular case where the boxes are connected. This is reminiscent of [16], where we also

475 showed that a certain property enjoyed by the λ -calculus can be extended to proof-structures,
476 provided they are connected inside boxes. We leave that work to a subsequent paper.

477 **Functoriality and naturality** Our functorial point of view on proof-structures might unify
478 many results. Let us cite two of them:

- 479 ■ a sequent calculus proof of $\vdash \Gamma$ can be translated into a path from the empty sequence
- 480 into Γ . This could be the starting point for the formulation of a new correctness criterion;
- 481 ■ the category **Path** can be extended with higher structure, allowing to represent cut-
- 482 elimination. The functors \mathbf{qMELL}^{\times} and $\mathfrak{PqDiLL}_0^{\times}$ can also be extended to such higher
- 483 functors, proving via naturality that cut-elimination and the Taylor expansion commute.

484 **Acknowledgments** This work has been partially funded by the EPSRC grant EP/R029121/1
485 “Typed Lambda-Calculi with Sharing and Unsharing” and the ANR project Rapido (ANR-
486 14-CE35-0007).

487 — References —

- 488 1 Denis B chet. Minimality of the correctness criterion for multiplicative proof nets. *Mathematical*
489 *Structures in Computer Science*, 8(6):543–558, 1998.
- 490 2 Dennis V. Borisov and Yuri I. Manin. *Generalized Operads and Their Inner Cohomomorphisms*,
491 volume 265 of *Progress in Mathematics*, pages 247–308. Birkh user Basel, Basel, 2007.
492 doi:10.1007/978-3-7643-8608-5_4.
- 493 3 Pierre Boudes. Thick subtrees, games and experiments. In *Typed Lambda Calculi and*
494 *Applications, 9th International Conference (TLCA 2009)*, volume 5608 of *Lecture Notes in*
495 *Computer Science*, pages 65–79. Springer, 2009. doi:10.1007/978-3-642-02273-9_7.
- 496 4 Pierre Boudes, Fanny He, and Michele Pagani. A characterization of the taylor expansion
497 of lambda-terms. In *Computer Science Logic 2013 (CSL 2013)*, volume 23 of *LIPICs*, pages
498 101–115. Schloss Dagstuhl - Leibniz-Zentrum fuer Informatik, 2013. doi:10.4230/LIPICs.
499 CSL.2013.101.
- 500 5 G rard Boudol. The lambda-calculus with multiplicities (abstract). In *4th International*
501 *Conference on Concurrency Theory (CONCUR '93)*, volume 715 of *Lecture Notes in Computer*
502 *Science*, pages 1–6. Springer, 1993. doi:10.1007/3-540-57208-2_1.
- 503 6 Daniel de Carvalho. The relational model is injective for multiplicative exponential linear
504 logic. In *25th Annual Conference on Computer Science Logic (CSL 2016)*, volume 62 of
505 *LIPICs*, pages 41:1–41:19. Schloss Dagstuhl - Leibniz-Zentrum fuer Informatik, 2016. doi:
506 10.4230/LIPICs.CSL.2016.41.
- 507 7 Daniel de Carvalho. Taylor expansion in linear logic is invertible. *Logical Methods in Computer*
508 *Science*, 14(4), 2018. doi:10.23638/LMCS-14(4:21)2018.
- 509 8 Daniel de Carvalho and Lorenzo Tortora de Falco. The relational model is injective for
510 multiplicative exponential linear logic (without weakenings). *Annals of Pure and Applied Logic*,
511 163(9):1210–1236, 2012. doi:10.1016/j.apal.2012.01.004.
- 512 9 Thomas Ehrhard. On K the sequence spaces and linear logic. *Mathematical Structures in*
513 *Computer Science*, 12(5):579–623, 2002. doi:10.1017/S0960129502003729.
- 514 10 Thomas Ehrhard. Finiteness spaces. *Mathematical Structures in Computer Science*, 15(4):615–
515 646, 2005. doi:10.1017/S0960129504004645.
- 516 11 Thomas Ehrhard and Laurent Regnier. The differential lambda-calculus. *Theoretical Computer*
517 *Science*, 309(1-3):1–41, 2003. doi:10.1016/S0304-3975(03)00392-X.
- 518 12 Thomas Ehrhard and Laurent Regnier. B hm Trees, Krivine’s Machine and the Taylor
519 Expansion of Lambda-Terms. In *Second Conference on Computability in Europe (CiE 2006)*,
520 volume 3988 of *Lecture Notes in Computer Science*, pages 186–197. Springer, 2006. doi:
521 10.1007/11780342_20.

- 522 13 Thomas Ehrhard and Laurent Regnier. Differential interaction nets. *Theoretical Computer*
523 *Science*, 364(2):166–195, 2006. doi:10.1016/j.tcs.2006.08.003.
- 524 14 Thomas Ehrhard and Laurent Regnier. Uniformity and the Taylor expansion of ordinary
525 lambda-terms. *Theoretical Computer Science*, 403(2-3):347–372, 2008. doi:10.1016/j.tcs.
526 2008.06.001.
- 527 15 Jean-Yves Girard. Linear logic. *Theoretical Computer Science*, 50(1):1–101, 1987. doi:
528 10.1016/0304-3975(87)90045-4.
- 529 16 Giulio Guerrieri, Luc Pellissier, and Lorenzo Tortora de Falco. Computing connected proof(-
530 structure)s from their Taylor expansion. In *1st International Conference on Formal Structures*
531 *for Computation and Deduction (FSCD 2016)*, volume 52 of *LIPICs*, pages 20:1–20:18. Schloss
532 Dagstuhl - Leibniz-Zentrum fuer Informatik, 2016. doi:10.4230/LIPICs.FSCD.2016.20.
- 533 17 Giulio Guerrieri, Luc Pellissier, and Lorenzo Tortora de Falco. Proof-net as graph, taylor
534 expansion as pullback. In *Logic, Language, Information, and Computation - 26th International*
535 *Workshop (WoLLIC 2019)*, volume 11541 of *Lecture Notes in Computer Science*, pages 282–300.
536 Springer, 2019. doi:10.1007/978-3-662-59533-6_18.
- 537 18 Emma Kerinec, Giulio Manzonetto, and Michele Pagani. Revisiting call-by-value Böhm trees
538 in light of their Taylor expansion. *CoRR*, abs/1809.02659, 2018. URL: <http://arxiv.org/abs/1809.02659>.
- 539 19 Giulio Manzonetto and Domenico Ruoppolo. Relational graph models, Taylor expansion
540 and extensionality. *Electronic Notes in Theoretical Compututer Science*, 308:245–272, 2014.
541 doi:10.1016/j.entcs.2014.10.014.
- 542 20 Damiano Mazza and Michele Pagani. The separation theorem for differential interaction
543 nets. In *Logic for Programming, Artificial Intelligence, and Reasoning, 14th International*
544 *Conference (LPAR 2007)*, volume 4790 of *Lecture Notes in Computer Science*, pages 393–407.
545 Springer, 2007. doi:10.1007/978-3-540-75560-9_29.
- 546 21 Damiano Mazza, Luc Pellissier, and Pierre Vial. Polyadic approximations, fibrations and
547 intersection types. *PACMPL*, 2(POPL):6:1–6:28, 2018. doi:10.1145/3158094.
- 548 22 Michele Pagani and Christine Tasson. The inverse taylor expansion problem in linear logic. In
549 *24th Annual Symposium on Logic in Computer Science (LICS 2009)*, pages 222–231. IEEE
550 Computer Society, 2009. doi:10.1109/LICS.2009.35.
- 551 23 Michele Pagani, Christine Tasson, and Lionel Vaux. Strong normalizability as a finite-
552 ness structure via the taylor expansion of λ -terms. In *Foundations of Soft-*
553 *ware Science and Computation Structures - 19th International Conference (FOSSACS*
554 *2016)*, volume 9634 of *Lecture Notes in Computer Science*, pages 408–423. Springer, 2016.
555 doi:10.1007/978-3-662-49630-5_24.
- 556 24 Christian Retoré. A semantic characterisation of the correctness of a proof net. *Mathematical*
557 *Structures in Computer Science*, 7(5):445–452, 1997. doi:10.1017/S096012959700234X.
- 558 25 Christine Tasson. *Sémantiques et Syntaxes Vectorielles de la Logique Linéaire*. PhD thesis,
559 Université Paris Diderot, France, December 2009. URL: <https://tel.archives-ouvertes.fr/tel-00440752>.
- 560 26 Lionel Vaux. Taylor expansion, lambda-reduction and normalization. In *26th EACSL Annual*
561 *Conference on Computer Science Logic (CSL 2017)*, volume 82 of *LIPICs*, pages 39:1–39:16.
562 Schloss Dagstuhl - Leibniz-Zentrum fuer Informatik, 2017. doi:10.4230/LIPICs.CSL.2017.39.